

The Hyperfine Structure of Antiprotonic Helium and the Antiproton Magnetic Moment

T. Pask¹, B. Juhasz¹, E. Widmann¹

¹*Stefan Meyer Institute for Subatomic Physics, Austrian Academy of Sciences,
Boltzmannngasse 3, 1090 Wien, Austria*

The ASACUSA collaboration is performing laser and microwave spectroscopy of antiprotonic helium ($\bar{p}\text{He}^+$), a metastable three-body system consisting of a helium nucleus, an antiproton and an electron, at the Antiproton Decelerator (AD) of CERN. $\bar{p}\text{He}^+$ exhibits a hyperfine splitting (HFS) which is unique due to the large angular momentum of the metastable states ($L_{\bar{p}} \approx 35$): the HFS consists of a dominant splitting caused by the interaction of $L_{\bar{p}}$ with the electron spin S_e and a smaller splitting due to the interaction of the antiproton magnetic moment $\mu_{\bar{p}}$ with the other moments. The hyperfine splitting has been measured for the first time in 2001 with a precision of 3×10^{-5} [1]. The two observed transitions are in agreement with QED calculations at a level of 6×10^{-5} , which corresponds to the theoretical accuracy. The agreement gives a limit on the antiproton *orbital* g -factor of $|g_l^{\bar{p}} - 1| < 6 \times 10^{-5}$ [2]. The difference of the two transition frequencies is directly related to the value of the *spin* magnetic moment $\mu_{\bar{p}}$, which so far is known to only 0.3%. ASACUSA has started a new measurement with the goal of increasing the experimental precision by an order of magnitude, which would lead to a determination of $\mu_{\bar{p}}$ to 0.1%.

References

- [1] E. Widmann *et al.*, Phys. Rev. Lett. **89**, 243402 (2002).
- [2] D. Bakalov and E. Widmann, Phys. Rev. A **76**, 012512 (2007).